

WARNEK LAMBERTI CO. LLC *EP 1348701-A1
2002.03.28 2002-290788(+2002EP-290788) (2003.10.01) C07D
277/42, A61K 31/426, C07D 417/04, A61K 31/427

New (2,4-disubstituted-thiazol-5-yl)amine compounds useful for treating diseases e.g. osteoarthritis, multiple sclerosis, osteoporosis, asthma, cancer and graft rejection (Eng)

C2003-235032 R(AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI
LT LU LV MC MK NL PT RO SE SI TR)

Addnl. Data: VERGNE F, BERNARDELLI P, LORTHIOIS E, DUCROT F


NOVELTY

(2,4-Disubstituted-thiazol-5-yl)amine compounds (I), their racemic forms, isomers, N-oxides, and their acidic or basic salt forms are new.

DETAILED DESCRIPTION

(2,4-Disubstituted-thiazol-5-yl)amine compounds of formula (I), their racemic forms, isomers, N-oxides, and their acidic or basic salt forms are new.

R_3-COOH
reactant



$R_{1a} = H$ or (aryl)1-6C alkyl;

$R_{1b} =$ (hetero)cycloalkyl, or (hetero)aryl (all optionally substituted by halo, trifluoromethyl, nitro, cyano, oxo, $-NR_4R_5$, $-CO_2R_4$, $-CONR_4R_5$, $-OR_4$, $-S(O)_mR_4$, $-S(O)_mR_4R_5$, tetrazolyl or 1-6C alkyl (optionally mono- or tri-substituted by $-OR_4$, $-NR_4R_5$ or $-CO_2R_4$));

n and $m = 0 - 2$;

R_4 and $R_5 = H$ or $-X_1R_6$;

X_1 and $X_2 =$ single bond or 1-6C alkylene;

EP 1348701-A+

$R_4 =$ 1-6C alkyl, (hetero)cycloalkyl or (hetero)aryl;
 $R_2 =$ 1-6C alkyl, 2-6C alkenyl, 2-6C alkynyl, aryl or cycloalkyl;
 $R_3 =$ (hetero)cycloalkyl or (hetero)aryl (all optionally substituted by halogen, nitro, cyano, trifluoromethyl, oxo, 1-6C alkyl, $-OR_6$, $-NR_6R_7$, $-COR_6$, $-CO_2R_6$, $-CONHOH$, $-CONR_6R_7$, $-S(O)_mR_6$, $-S(O)_m-NR_6R_7$, $-NR_6COR_7$, $-NR_6SO_2R_7$, $-N(SO_2R_7)_2$, $-NR_6-CO-NR_7R_8$ or tetrazolyl);

R_6 and $R_7 = H$ or $-X_2R_6$;

$R_8 =$ 1-6C alkyl, (hetero)cycloalkyl or (hetero)aryl (all optionally mono- or tri-substituted by OH, 1-6C alkoxy, 1-6C alkyl, amino, mono-1-6C alkylamino, di-1-6C alkylamino, carboxy, 1-6C alkoxy carbonyl or benzyl); and

$R_8 = H$ or 1-6C alkyl.

The aryl is an aromatic monocyclic or bicyclic system containing 5-10C; in the bicyclic ring system, one of the rings is aromatic and the other ring is optionally aromatic or partially hydrogenated and when the second ring is partially hydrogenated, then the ring is optionally mono- or di-substituted by oxo. The heteroaryl is the aryl group in which 1-4 carbon atoms are replaced by 1-4 heteroatoms selected from O, S and N. The cycloalkyl is a monocyclic or polycyclic system containing 3-10C and is saturated or partially unsaturated but without aromatic character and in the polycyclic system, each cycle could be

fused together or formed a link. The heterocycloalkyl is the cycloalkyl group in which 1-4 carbon atoms are replaced by 1-4 heteroatoms selected from O, S or N.

An INDEPENDENT CLAIM is included for preparation of (I).

ACTIVITY

Immunosuppressive; Antiinflammatory; Respiratory-Gen.; CNS-Gen.; Antiallergic; Gastrointestinal-Gen.; Analgesic; Osteopathic; Neuroprotective; Antiasthmatic; Cytostatic; Anti-HIV; Antiarthritic.

MECHANISM OF ACTION

Phosphodiesterase-7 (PDE-7) inhibitor.

(I) were tested for inhibition of cyclic nucleotide phosphodiesterase 7, as given in W.J. Thompson et al. 1979, Advances in Cyclic Nucleotide Research, Vol.10:69-92, ed.G. Brooker et al. Raven Press, NY. They showed IC_{50} value of 0.02 - 100 micro M. No results for specific compounds are given.

USE

For the treatment of a disease including T-cell related disease, autoimmune disease, inflammatory disease, respiratory disease, CNS

EP 1348701-A+/1

2003-835650/78

disease, allergic diseases, endocrine or exocrine pancreas disease, gastrointestinal diseases, visceral pain, inflammatory bowel disease, osteoarthritis, multiple sclerosis, osteoporosis, chronic obstructive pulmonary disease (COPD), allergic rhinitis, asthma, cancer, acquired immune deficiency syndrome (AIDS) and graft rejection (claimed).


ADVANTAGE

The compounds are active at very low concentrations.

SPECIFIC COMPOUNDS

4 Compounds (I) are specifically claimed, e.g. 7-[5-(cyclohexylamino)-4-methyl-1,3-thiazol-2-yl]quinazoline-4-amine (Ia).

BEST AVAILABLE COPY



ADMINISTRATION

The compounds, in a dosage of 1 mg - 1 g per day, can be administered orally, parenterally (including intravenously, intramuscularly or subcutaneously), per- or trans-cutaneously, intravaginally, rectally, nasally, perlingually, buccally, ocularly or by respiratory route.

EXAMPLE

To a solution of 4-oxo-3,4-dihydro-quinazoline-7-carboxylic acid in tetrahydrofuran (THF) was added 1,1'-carbonyldiimidazole (I.2

EP 1348701-A+/2

(con't)

(C) 2004 Copyright Derweni Information Ltd.

equivalents) and the mixture was stirred for 30 minutes. The (S)-L-alanine tert-butyl ester was added and the mixture was stirred for 24 hours. The solvent was removed and the residue was worked up to obtain tert-butyl(2S)-2-[[[(4-oxo-3,4-dihydroquinazolin-7-yl)carbonyl]amino]propanoate (A).

(A) was added to a solution of 5% trifluoroacetic acid in dichloromethane and the mixture was stirred for 3 hours, followed by a work-up to obtain (2S)-2-[[[(4-oxo-3,4-dihydroquinazolin-7-yl)carbonyl]amino]propanoic acid (A1).

To a solution of (A1) in THF, 1,1'-carbonyldiimidazole (1.2 equivalents) was added and the mixture was stirred for 30 minutes. Then the cyclohexylamine was added and the mixture was stirred for 24 hours, followed by a work-up to obtain N-[(1S)-2-(cyclohexylamino)-1-methyl-2-oxoethyl]-4-oxo-3,4-dihydroquinazoline-7-carboxamide (A2).

To a solution of (A2) in pyridine was added Lawesson's reagent and the mixture was heated to 100°C for 6 hours. After cooling to room temperature, saturated aqueous sodium hydrogen carbonate solution was added to the reaction mixture and the product was worked up to obtain 7-[5-(cyclohexylamino)-4-methyl-1,3-thiazol-2-yl]quinazoline-4(3H)thione (A3).

To a stirring solution of (A3) and potassium carbonate (1.2

equivalents) in methanol was added CH₃I. After 30 minutes, the solvent was removed to obtain N-cyclohexyl-4-methyl-2-[4-(methylthio)quinazolin-7-yl]-1,3-thiazol-5-amine (A4).

A solution of (A4) in saturated butanolic ammonia was sealed in steel bomb and heated at 100°C for 2 days. The solvent was removed and the residue was purified to obtain 7-[5-(cyclohexylamino)-4-methyl-1,3-thiazol-2-yl]quinazoline-4-amine (Ia).

DEFINITIONS

Preferred Definitions:

R_{1a} = H;

R_{1b} = cyclohexyl group (optionally mono-substituted by OH) or phenyl (optionally mono-substituted by tetrazolyl or -CO₂R₄);

R₄ = H or 1-6C alkyl;

R₂ = methyl;

R₃ = quinoxalinylnyl, 1H-quinazolinylnyl, 3H-quinazolinylnyl-4-one, 1H-quinazolinylnyl-2,4-dione (all optionally mono- tri-substituted by halo, 1-6C alkyl, OR₆ or NR₆R₇);

X₂ = single bond;

R₆ = 1-6C alkyl (optionally monosubstituted by OH, 1-6C alkoxy, amino, mono-1-6C alkylamino or di-1-6C alkylamino).

EP 1348701-A+3

2003-835650/78

TECHNOLOGY FOCUS

Organic Chemistry - Preparation (Claimed): Preparation of (I) involves:

- (1) coupling a carboxylic acid of formula R₃-C(=O)OH with an amine of formula ProtO-C(=O)-CHR₂-NH₂ under peptidic coupling conditions to give a coupled product of formula R₃-C(=O)-NH-CH(CO₂Prot)-R₂;
- (2) deprotecting the coupled product by treatment with an acid or base to give a free carboxylic acid compound of formula R₃-C(=O)-NH-CH(CO₂H)-R₂ (V);
- (3) reacting (V) with a primary amine of formula R_{1b}-NH₂ under peptidic coupling conditions in the presence of a coupling agent to give a couple product of formula R₃-C(=O)-NH-CH(R₂)-C(=O)-NH(R_{1b}) (VI);
- (4) treating (VI) with Lawesson's reagent in basic medium to give (I) (in which R_{1a} is H);
- (5) treating (I) (in which R_{1a} is H) with R'_{1a}-L₁ to give (I) (in which R_{1a} is R'_{1a}) optionally under alkaline medium;
- (6) purifying (I) (in which R_{1a} is H or R'_{1a}) by a conventional purifying

technique; and

- (7) separating into their respective isomers, followed by converting into their salts with acid or base or into N-oxide.

Prot = protective group of carboxylic acid group;

R'_{1a} = (aryl)1-6C alkyl;

L₁ = leaving group.

(20pp8014DwgNo.0/0)

EP 1348701-A/4